

Temporal scale of field experiments in benthic ecology

Jenkins, Stuart; Yrayzoz, Marc Uya

Marine Ecology: progress series

DOI:

[10.3354/meps11659](https://doi.org/10.3354/meps11659)

Published: 01/04/2016

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Jenkins, S., & Yrayzoz, M. U. (2016). Temporal scale of field experiments in benthic ecology. *Marine Ecology: progress series*, 547, 273-286. <https://doi.org/10.3354/meps11659>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1
2
3
4
5
6
7
8
9

Temporal scale of field experiments in benthic ecology

Stuart R Jenkins and Marc Uyà

School of Ocean Sciences, Bangor University, Menai Bridge, Anglesey, UK LL59 5AB

Running head: Duration of experiments in benthic ecology

Email: s.jenkins@bangor.ac.uk

Abstract

The issue of scale in ecology is all-pervasive, with recognition that most ecological questions are scale dependent. Scaling up has become part of the lexicon of experimental ecology, and in marine benthic systems has resulted in numerous advances. While it is clear that manipulative experiments in benthic systems have been conducted over increasing spatial scales, it is less clear whether the notion of scaling up has been applied to temporal scale. Here, we examine the temporal scale at which experiments have been undertaken before reviewing longer term studies to examine the insights gained from extending the duration of observation following perturbation. Field experiments which examined population/community responses to perturbations, and studies which monitored the consequences of natural disturbances, were identified over the period 1980-2013. The median length of study was 10 months and only 12% of studies were carried out over more than 3 years. Neither the median study length, nor the proportion of studies longer than 2 or 3 years, showed a trend over the 33 years. Review of experiments with a duration of 3 years or more revealed numerous benefits of a long-term approach. Some of these were unexpected, but others were predictable based on life history traits of dominant organisms, slow successional patterns, or response variables related to longer term community level responses, such as stability. The review suggests modest investment in resources to extend the duration of experiments can bring substantial benefits and hence consideration of experimental duration should be one of the primary decisions in planning field experiments.

Keywords: field experiments; long term; perturbation; scaling up; temporal scale

Introduction

Our understanding of ecological processes in the marine benthic environment has been built on a foundation of manipulative experiments over the past half century or so, in a range of intertidal and shallow sub-tidal systems. Both soft and hard substrata habitats have been utilised by experimental ecologists, intent on understanding the processes structuring marine assemblages, and hence explaining patterns in abundance, distribution and diversity of species (see Raffaelli & Hawkins 1996 for review). Marine ecological experiments have also contributed to important advances in ecological theory (e.g. Connell 1961, Paine 1966, Dayton 1975, Menge & Sutherland 1976, Bertness & Callaway 1994). The intertidal zone is favoured by experimental ecologists owing to the logistical ease of working on marine communities when the tide is out, combined with a sharp marine/terrestrial environmental gradient which is conducive to investigations of the interaction of physical and biological processes. Additionally, the intertidal is inhabited by organisms which are generally small and sessile/slow moving and hence amenable to experimental manipulation (Connell 1972, Paine 1977, Underwood 2000) and in the case of hard substrate, the environment is two-dimensional in nature. Sub-tidal work is more challenging, but there is still a vast body of experimental research undertaken in this environment, either through the use of SCUBA or suspending experiments from floating or fixed structures. Focal habitats for experimental research in the coastal zone include coral reefs, seagrass beds, subtidal rocky reefs, intertidal rocky shores, mud-flats and beaches and sub-tidal soft sediment systems.

The development of experimental ecology in marine systems and the advances made over the past 50 years or so are well documented (see Paine 1994, Underwood 2000, Castilla 2000 for reviews). One area which has received increasing attention, in marine as well as terrestrial and freshwater habitats, is the issue of scale (Wiens 1989, Levin 1992, Schneider 2001, Ellis & Schneider 2008). Schneider (2001) neatly summarised three components to the problem of scale: problems in ecology often exist at decadal time scales over large ecosystems; most variables can only be measured in small areas over short time scales; patterns and processes observed at small scales do not necessarily hold at large scales. It is increasingly recognised that environmental drivers and ecological processes operate over a range of different spatial and temporal scales and over the past two decades, scale, in a quantitative sense, has been increasingly incorporated into experimental work (Ellis & Schneider 2008).

Ecologists in coastal marine systems have been key in developing a framework in which understanding from manipulative experiments may be scaled up through the comparative experimental approach (Menge et al. 2002). Identical replicated experiments are conducted at multiple sites often over some environmental gradient (e.g. Menge et al. 1997, Jenkins et al. 2001, Coleman et al. 2006), thus allowing determination of the extent to which large scale physical conditions, which are not amenable to manipulation, can modify ecological processes. This 'scaling up' approach has led to a number of important developments in understanding of coastal systems, particularly in recognising the importance of latitudinal variation in temperature (e.g. Leonard 1999, Sanford 1999, Jenkins et al. 2001), regional variation in productivity (Menge et al. 1997, Broitman et al. 2001) and large scale variation in recruitment regime (Connolly & Roughgarden 1998). Studies of systems in contrasting oceanographic regimes have made progress in integrating small, medium and large scale processes and in particular have shown that benthic assemblages in shallow water or intertidal areas do not operate in isolation from the planktonic, pelagic realm (Menge et al. 2003).

In considering scale, attention has not solely focused on variation in space, with increasingly sophisticated ecological experiments examining processes at a range of different temporal scales. For example the importance of seasonality in determining the magnitude and direction of ecological processes is now recognised in experiments through manipulations being performed at different times of the year, and it is not uncommon for multiple starting dates to be nested within season in order to provide unconfounded tests of seasonal effects (e.g. Jenkins et al. 2005). Another aspect of temporal scale is the variability of events through time. Benedetti-Cecchi (2003) highlighted that temporal variability in the occurrence of perturbations is generally expressed in terms of the frequency of events, but argues that a more appropriate approach which avoids confounding the variance with the mean effect size (i.e. a more frequent disturbance is also more intense) is to design experiments in which levels of intensity, and those of variability, are chosen independently. His work has led to numerous studies explicitly assessing the importance of temporal variance in perturbations (e.g. Incera et al. 2010, Maggi et al. 2012, Oliveira et al. 2014).

Consideration of temporal scale also clearly needs to take account of the duration of any particular manipulative experiment. Underwood (2000), based on the assertion of Connell & Sousa (1983) that studies should encompass a period as long as the turnover of individuals in resident species, suggests that on Australian rocky shores a period of between 5 and 7 years might be appropriate, but also considers studies may need to be much longer to allow analyses

of equilibria. Yodzis (1988) considered that for the impact of a press perturbation to be known, a long term approach is required. He recommended adopting a rule of thumb such that the time scale of observation should approximate to two times the sum of the generation times of a pressed species (i.e. one whose density has been modified) and the potentially impacted species of interest. It is debateable whether all experiments require such duration, and clearly there are specific questions which can be addressed by much shorter studies. For example Menge (1997) tested the assertion that indirect effects may take longer to become apparent, following a manipulation, than direct effects, by determining the relationship between duration of experiment and the importance of indirect effects using 23 rocky intertidal interaction webs. He concluded that in the case studies examined, which ranged in length from 0.5 to 156 months, '*interpretations of community dynamics are not likely to be distorted because experiments were too brief*' (p.812). Thus, based on addressing a very specific goal, that of quantifying indirect effects in community dynamics, Menge (1997) considered the experiments he reviewed as being sufficiently long.

In most fields of marine benthic ecology there has been a lack of explicit consideration of experimental duration. A field of burgeoning research that has recently recognised the serious implications of a short term approach is in Biodiversity and Ecosystem Functioning (BEF) work (Hooper et al. 2005, O'Connor & Crowe 2005, Raffaelli 2006, Crowe et al. 2012, Duffy et al. 2012). Crowe et al. (2012) used existing meta-analyses (Cardinale et al. 2007, Stachowicz et al. 2007) to review the temporal scale of biodiversity-ecosystem functioning (BEF) studies conducted in both laboratory and field settings (in marine, terrestrial and freshwater environments). The authors argue that limiting the temporal extent of experiments has a marked effect on the conclusions regarding the importance of biodiversity in ecosystems and the mechanism by which biodiversity may exert an effect; short term experiments will over-emphasise the importance of dominant species (a species identity effect) relative to complementarity among species. Stachowicz et al. (2008a) examined the effect of seaweed diversity on cover over a period of three years. They showed that seaweed cover in diverse treatments only exceeded that in the best performing monoculture after a period of 18 months, a period much longer than many biodiversity studies (Crowe et al. 2012). These observations might reasonably be seen as strong arguments for lengthening the duration of BEF experiments (although doing so in any multi-trophic manner incorporating large, mobile long lived consumers will be enormously challenging (Raffaelli 2006)).

Although many aspects of temporal scale have been addressed through ecological experiments in recent decades, it is debateable whether the scaling up approach, so prevalent in considerations of spatial scale, has been applied to temporal scale. Thus it is unclear whether the overall length of experimental study (i.e. scaling up in time) has increased, despite numerous publications tentatively expressing concern at the paucity of longer term experimentation (e.g. Connell & Sousa 1983, Underwood 2000, Raffaelli 2006). Such calls have no doubt been influenced by the perceived benefits of long term experimentation (Underwood 2000; Crowe et al 2012). Here, we examine the temporal scale at which ecological experiments (which apply some form of perturbation) are undertaken in intertidal and shallow sub-littoral environments before reviewing the literature to highlight the range of insights gained from long term experiments. We finish by considering the challenges associated with increasing the temporal extent of marine benthic experiments and the way in which these may be addressed.

Review of experimental duration

As outlined above there have been a number of calls for longer term experimentation, but a factor which may act counter to such a move is the increasing pressure on scientists to publish their work. While publication of research in a vibrant international literature is the life-blood of science, the pressure to publish could be argued to lead to a short term approach, with researchers rushing to publish each new piece of research. It is very easy to see how this short term approach can influence our approach to experimental ecology. Why spend 5 years undertaking an experiment when a paper can be published on the basis of results collected over a period of 18 months? By reviewing the literature (including both marine focused and general ecological journals) we hope to determine what range of temporal scales experimental marine ecologists are working at and whether the approach to experimental duration has changed over a 33 year period. This review is limited to assessment of the effects of perturbation on marine systems; thus long term monitoring *per se* is not considered although it is acknowledged that this approach may often address similar questions.

Methods

We examined literature published over a 33 year period in two of the leading marine ecological journals, Marine Ecology Progress Series (MEPS) and Journal of Experimental Marine Biology and Ecology (JEMBE), and three general ecological journals, Ecology, Oikos and Oecologia in order to examine the temporal scale and sampling frequency of field

perturbation experiments in shallow sub-tidal and intertidal benthic environments. In selecting publications, all types of pulse and press perturbations were included, from manipulations which simulate some form of natural or anthropogenic disturbance event, to those in which the density or presence of one or more particular species is altered. In addition to manipulation of biota we included perturbations to the physical environment, for example modification of the nutrient or light regime. One very common type of study was one in which new substrata (often simple PVC panels) were introduced into the environment. Such perturbations may be considered analogous to clearance of biota in natural communities, since both provide bare space, free from competitors.

In considering which studies to include, the main criterion beyond inclusion of an appropriate perturbation was the response variable measured. In general, studies were included which measured the community or population response of macro-organisms to a perturbation (i.e. meiofauna and microbial communities were rejected). Studies where the research aim was solely to examine some process rate, such as growth or mortality, or the effect of disturbance on an organisms' body condition, were excluded. In addition, studies which examined only a behavioural response were not included. One difficult judgement to be made was in studies designed to examine spatial or temporal patterns in the arrival of animals or algae i.e. settlement/recruitment. Where the aims of such work were only short term in nature, i.e. examination of settlement, such studies were not included since settlement can be clearly defined as a short term process (Jenkins et al. 2009). However, use of the term recruitment is more problematic. Because of its variable use in benthic ecology with no defined time period over which it applies (Keough & Downes 1982, Jenkins et al. 2009) studies examining recruitment were included unless the aim of the study was clearly related to understanding processes determining the arrival of propagules and subsequent settlement (e.g. Reinhardt et al. 2013) rather than the development of a benthic population or community.

In addition to manipulative experiments implemented by the experimenter, numerous studies examine the consequences of events or implementation of management structures not under the control of the experimenter. These include investigations of natural disturbances such as extreme weather events (e.g. Tegner et al. 1997) or human-induced perturbations such as pollution (e.g. Jacobs 1980), the exclusion of humans from Marine Protected Areas (Castilla & Bustamante 1989) and managed realignment of coastal limits through intentional flooding (Hughes et al. 2009). These 'natural experiments' were included in our review of literature

whenever data were available prior to the perturbation taking place, such that a true temporal control existed.

Using the criteria outlined above, all publications in MEPS, JEMBE, Ecology, Oikos and Oecologia at four yearly intervals from 1981 to 2013 were initially reviewed by examination of the title and abstract. Following identification of potential papers the full text was read and then either rejected or the appropriate information extracted. This approach was taken rather than performing a search based on keywords because of the difficulty in creating suitable search terms which would highlight appropriate papers without bias. Our search through 12,557 titles (6940 in general ecological journals; 5617 in marine ecological journals) identified a total of 263 publications containing 322 independent experiments over the nine publication years. These represented an average of 3.5% and 1.3% of the studies published in marine and general ecological journals respectively (or 15.9% of the marine related papers published in general journals). Although there was no obvious directional change over the 33 years of review in the proportion of appropriate studies selected in each year in either journal type, there was a general increase in the number of studies from which data were extracted owing to the huge increase in number of publications in all journals over time (from 299/564 in 1981 to 823/831 in 2013 for marine/general ecological journals respectively). In order to allow good representation of experimental work in earlier years, additional publications were sought for the earliest five dates by examining studies published up to two years before (and in the case of 1981, two years after). This approach was taken in the same way as described above, until a minimum of 10 independent experiments in both marine ecological and general ecological journals were identified for each of the four-yearly periods. This resulted in 293 publications containing 352 studies being reviewed.

The temporal scale of experiments in all 352 studies was determined as the length of time from initial sampling, immediately prior to the perturbation, to the final sampling event. Where multiple experiments within a single publication were conducted with the aim to examine a single goal or hypothesis, the maximum length of experimentation was documented. In determining the sampling interval (i.e. length of time between each sampling event) an average value over the period of the study was calculated. For example, in a ten month study examining the influence of bivalve shell material introduced to a mudflat on colonisation by burrowing thalassinidean shrimp, Feldman et al. (1997) made observations 1 month after the perturbation and after ten months. Here, the average sampling interval was recorded as 5 months. The habitats in which experiments were undertaken were classified as

inter-tidal or sub-tidal. In micro-tidal regimes such as the Mediterranean (e.g. Maggi et al. 2009) and Baltic (e.g. Lindegarth et al. 2001) studies undertaken in areas frequently exposed to air through variations in pressure and wind were classified as intertidal. The Caribbean fringing reef studied by Kilar & MacLachlan (1989) is exposed to air only during particular conditions, at certain times of the year, and was therefore considered sub-tidal. The classification of habitat type (rock, soft sediment, sea-grass, coral, mangrove, salt-marsh, artificial substrate) was generally clear cut. Cobble and boulder habitat were considered simply as rock rather than creating a novel category.

Classification of experiments as pulse or press was clear-cut in most cases. Pulse experiments were defined as those receiving a one-off, perturbation which was generally short in temporal extent compared to the temporal scale of the study. The introduction of artificial substrata such as PVC panels (e.g. Valdivia et al. 2005) or newly quarried boulders (e.g. Chapman 2013) was considered a pulse disturbance, analogous in many ways to the creation of bare space by a single disturbance event. In cases where both pulse and press perturbations were implemented, the experiments were invariably classified as press. For example in numerous studies, novel substrata were introduced in order to examine the consequences of a particular press manipulation, such as light and sediment (Connell 2005) or herbivores (Kaehler & Williams 1997). Other studies (e.g. Russell & Connell 2005) combined pulse manipulations such as the one-off removal of a particular species with continued press perturbations such as addition of nutrients. Rather than class these as a combination of both pulse and press, they were classed as press. In some studies, multiple perturbations were applied through time (Jewett et al. 2005, Incera et al. 2009, Minchinton & Fels 2013) and again here it was considered that the principle of an ongoing perturbation (although not continuous) justified categorisation as press.

Results and Discussion

64 % of the 352 studies over the 33 years were from marine journals (MEPS and JEMBE) whereas 36 % were published in general ecological journals (OIKOS, Ecology, Oecologia). The studies were split almost exactly equally between the intertidal (49.7%) and the sub-tidal (50.3%) zones. Of the eleven different habitats identified the most common was inter-tidal rocky shore (31% of studies) followed by sub-tidal rocky reef (14%) and then sub-tidal artificial substrate and sub-tidal coral reef (9% each). 62 % of the studies were categorised as press (217 studies) whilst the rest were considered as pulse (135).

Examination of the frequency distribution of study length across 33 years showed a clear skew to a shorter time frame (Figure 1). The highest proportion (24% and 33% for marine and general ecological journals respectively) were very short (between 0 and 3 months), with obvious peaks at 9-12 and 21-24 months for marine journals (presumably due to the logical attraction of terminating experiments at one and two years respectively) (Figure 1). Most studies (85% for marine and 82% for general ecological journals) lasted two years or less and only 39 of the 352 studies were conducted over a period of 3 years or more. Of these, 17 studies were of a period of 5 years or more. The median length of study over the period reviewed was 11 months in marine journals, 9 months in general ecological journals and 10 months overall (Figure 2). Although the median study length varied among years for marine journals (Kruskal Wallis test, Chi-Square = 19.427; df = 8; P = 0.013) there was no apparent trend in study length with time.

Study length showed differences between habitat (intertidal versus subtidal) and between perturbation type (pulse versus press) (Figure 3). The median length of inter-tidal studies (12 months) was significantly longer than sub-tidal studies (8 months) (Mann Whitney U test: $Z = -2.407$; $p = 0.016$ (Figure 3A). In the intertidal zone, 30% of studies were conducted over 2 years or more, approximately double that in the sub-tidal. The median study duration of pulse experiments (14 months) was significantly longer than press experiments (6 months) (Mann Whitney U test: $Z = -4.762$; $p < 0.001$) (Figure 3A) and a far greater proportion lasted two years or longer (35% of pulse compared to only 16% for press).

There was a significant positive relationship between experimental duration and sampling interval ($F_{1,347} = 108.5$; $P < 0.001$). Thus, as experiments got longer the time between sampling events increased. However, the relationship was weak ($r^2 = 0.236$), driven partially by the fact that a sampling interval (the dependent variable) cannot be greater than the experimental duration (the predictor variable) and indeed many studies well beyond the median duration had short sampling intervals of between 0.5 and 3 months.

In discussing the need for a longer term approach Underwood (2000, p.68) states that '*the majority of ecological studies are quite short, usually three to five years*'. In fact, our analysis of 352 studies over 33 years shows that for experiments (i.e. where some form of manipulation or perturbation has taken place, as opposed to purely observational work) the median length of study was only 10 months and only 12% of published work was over a period greater than 3 years. Patterns were similar irrespective of the type of journal (marine

specific versus general ecological) in which papers were published. These results were quite clear in indicating that calls for an increase in the temporal scale of studies have not led to any general increase in experimental duration over the past 3 decades. This is true whether considering the median duration of studies or the proportion which are longer than 24 or 36 months.

Different approaches can be taken to make an argument for the benefits of longer term experimental duration. One is in demonstrating that longer term experiments reveal, for example, differences in effect sizes compared to short term experiments. Such an approach is fraught with problems in designing a non-biased analysis, since often the benefits of long term experimentation are revealed not in a single bigger outcome but in multi-faceted results which provide greater insight into ecological complexities. Below we address such complexity by reviewing a range of longer term studies (generally greater than three years) which have revealed greater insight than could be achieved by equivalent short term work.

Insights from long term experiments

Controlled manipulations

There are a diversity of reasons for conducting long term experiments including: study of long lived species (e.g. Clarke & Allaway 1993; Jenkins et al. 2004) and slow successional change (Yakovis et al 2005); work in polar regions where development rates are much slower (Konar 2007, Beuchel & Gulliksen 2008); the search for long lived ecological phenomena such as alternate stable states (Petraitis & Dudgeon 2005); determination of the nature and extent of temporal community fluctuations in disturbed systems (Dye 1998). Defining how long to run field experiments is a difficult task but the number of studies in marine coastal systems which show differences between control and treatment plots more than 5 years (Clarke & Allaway 1993, Cervin et al. 2005, Viejo 2009, Schiel & Lilley 2011) and in some cases more than 10 years (Dye 1998, Jenkins et al. 2004, Beuchel & Gulliksen 2008) after a perturbation suggest that to address many community level questions experiments should run for years rather than months. Differences among treatments are not simply limited to community structure but also to overall community productivity (Tait & Schiel 2011) and to community stability (Dye 1998). For example, Tait & Schiel (2011) showed that 90 months after removal of the fucoid canopy alga *Hormosira banksii*, community primary productivity was only between 40% to 60% that of control areas. The reasons for slow recovery may be because successional development is inhibited (sensu Connell & Slatyer 1977) by early colonisers (e.g. Sutherland

& Karlson 1977), communities are dominated by slow growing species (e.g. Jenkins et al. 2004) or the environment is conducive to slow development, for example in polar regions (e.g. Beuchel & Gulliksen 2008).

In polar regions disturbance can have a profound effect on community structure owing to very slow, and in many cases, an almost complete lack of recovery (Dayton & Oliver 1977, Barnes 1996, Konar 2007). For example, Konar (2007) found that 4 years after clearance of sub-tidal boulders on the Alaskan north coast, cover of biota had only reached 2%, despite the fact that bare surface in the community in general was scarce (< 10%). A similar lack of colonisation (a total of 2 serpulid polychaetes present after 3 years, and reports of bare surfaces after 5 years) on a settlement panel array was observed in McMurdo Sound, Antarctica (Dayton 1989). This was followed at some point by bryozoan, hydroid, soft coral, and sponge colonisation leading to 'heavily covered' surfaces (Dayton 1989, p.1486). Low recruitment and slow growth may characterise polar regions, with physically driven episodic events driving a punctuated succession (Dayton 1989). Such an environment clearly requires long term experimental work.

One argument for conducting long term experiments is that they can reveal quite unexpected results. Jenkins et al. (1999a,b, 2004) report the results of factorial manipulations of the long lived canopy species *Ascophyllum nodosum* and the keystone grazer *Patella vulgata* on sheltered canopy dominated shores of the Isle of Man. Short term results over a period of one to two years included the die back of understory turfing algae, opening up of bare space and replacement of the *Ascophyllum* canopy with alternative more opportunistic shorter lived fucoid species, *Fucus vesiculosus* and *F.serratus*. However it was the longer term observations over 6 years (Jenkins et al. 1999a) and then twelve years (Jenkins et al. 2004) which revealed the importance of *Ascophyllum* in limiting patellid limpet populations; loss of the facilitatory function of the canopy on understory turfing algae, and hence the opening up of bare space, resulted in enhancement of limpet recruitment, leading to populations up to six times greater in abundance over a decade after canopy loss. Here, experimental work demonstrated the stabilising function of turfing algae in the undisturbed community. Other work has recognised how the dense low lying nature of turfing algae, combined with sediment trapping ability (Connell et al. 2014) contributes to stability or slow succession (e.g. Cervin et al. 2005, Schiel & Lilley 2011).

It is frequently stated that coastal marine systems are ideal environments in which to conduct manipulative experiments because dominant organisms have relatively short generation times and hence experiments may be conducted over short time scales. However it is debatable whether generation times in primary producers, key consumers and dominant space occupiers (which are often a number of years) are short enough to justify this view (Beuchel & Gulliksen 2008). Species life history traits are important determinants for the recovery of communities following disturbance in a range of different ecosystems including saltmarshes (e.g. Bertness 1991), mangroves (e.g. Clarke & Allaway 1993) and rocky shores (e.g. Methratta & Petraitis 2008), and species longevity, as well as mode and rate of colonisation, have clear implications for experimental duration. Methratta & Petraitis (2008) examined recovery from disturbance on rocky shores focusing on species with contrasting longevity, barnacles and furoid algae. They showed that differences in life history traits of dominant species modified successional trajectories at different spatial scales over a 5 year period. Understanding of the population dynamics and community structuring role of long lived species will invariably require longer term studies than investigation into species with short generation times, clearly demonstrated by the nine year study on mangrove regeneration of Clarke & Allaway (1993) and the three year study of Bertness (1991) on *Spartina* competitive interactions. On rocky shores *Ascophyllum nodosum*, a dioecious, long-lived species (Aberg 1992), with low rates of recruitment rate and slow growth rate (Aberg & Pavia 1997, Cervin et al. 2005) has been the focus of a number of long term studies in the NE and NW Atlantic (e.g. Jenkins et al. 1999a,b, 2004, Cervin et al. 2005, Ingolfsson & Hawkins 2008, Araújo et al. 2012). Araújo et al. (2012) showed that the impacts of human trampling can have long term (5 years) effects on understorey community structure in stands of *Ascophyllum nodosum*. Even small scale (25 × 25cm) disturbance events can have long lasting effects on *Ascophyllum* understorey communities (Cervin et al. 2005); small scale loss of *Ascophyllum* canopy led to a change in community structure still apparent after 7 years. Such effects can occur despite recovery of the long lived biomass dominants. Schiel & Lilley (2011) showed differences between disturbed and control understory communities 8 years after removal of the *Hormosira* canopy even though this dominant furoid returned to a nearly closed canopy. Communities composed of long lived individuals are often described as stable, but in reality may only be stable as a consequence of the long lived nature of their components (Connell & Sousa 1983). Long term observations following perturbations, especially where dominant biota are long lived, are essential to judge the stability of communities and consequent

properties of ecosystems (Farrell 1988, Hooper et al. 2005). Hooper et al. (2005) points out that theoretical work on stability has outpaced the ability of empirical work to test predictions, essentially because of the time and resource requirements of long term experimental work required to determine the fate of all adults over at least one complete turnover (Connell & Sousa 1983). The problem of time scale is one which has dogged the sometimes vigorous debate surrounding the existence, or otherwise, of alternate stable states (see Petraitis & Dudgeon 2004 for review). The question of whether systems can occupy more than one equilibrium point and hence support different stable communities in the same habitat (Lewontin 1969, Connell & Sousa 1983, Peterson 1984, Petraitis & Latham 1999, Petraitis et al. 2009) has been addressed in a number of different marine coastal assemblages including coral reefs (Scheffer et al. 2001), rocky shores (Paine et al. 1985, Petraitis & Latham 1999), sub-tidal rocky reefs (Simenstad et al. 1978) and soft sediments (van de Koppel et al. 2001). Part of the problem in answering what at first sight seems a relatively simple question is that spatial and temporal scale are an intractable part of defining stability. Connell & Sousa (1983) argued that experimental tests of stability require long term experiments where *'the fate of all adults of the population or community must either be followed for a minimal period of at least one complete turnover, or their replacement probabilities estimated'* (p.808). Others disagree (Peterson 1984, Sutherland 1990) and over the last decade or so Petraitis and co-workers (e.g. Petraitis & Latham 1999, Petraitis & Dudgeon 2005, Petraitis et al. 2009) have implemented manipulations which although long term by the standards of this review do not reach the temporal standards defined by Connell & Sousa (1983). The possibility that mussel beds and stands of *Ascophyllum nodosum* represent alternative states in sheltered bays of the Gulf of Maine was tested through initiating pulse disturbances (clearance of the *Ascophyllum* canopy) of different sizes and monitoring successional pathways over a nine year period (Petraitis et al. 2009). The shift from *Ascophyllum* assemblage to mussel bed in 37% of large clearances, nine years after disturbance, is argued by Petraitis et al. (2009) to show that these communities represent multiple stable states on the intertidal shore of the Gulf of Maine. They go on to present a conceptual framework which combines this view with the prevailing paradigm of environmentally driven consumer control (Lubchenco & Menge 1978, Bertness et al. 2002). Irrespective of tests of ecological theory, investigation of how the scale of disturbance affects successional trajectory clearly requires a long term approach in this environment at least, as evidenced by the diversity of successional responses among clearing sizes and sites over a six year period (Petraitis & Dudgeon 2005).

425 Natural perturbations and Marine Protected Areas

426 The majority of ecological experimentation involves small scale, controlled manipulation,
427 which, if of sufficient duration, may give insight into the time scale of recovery of marine
428 communities following perturbation. However, Schiel and Lilley (2011) point out the
429 limitation of such approaches in understanding recovery at coast-wide scales, where
430 propagule supply may be limiting. Underwood (1998, p.292) makes a cogent argument to
431 '*grab opportunities afforded by dramatic events, such as unusually severe weather and large-*
432 *scale changes brought about by natural agents*' to enable a longer term and larger scale
433 approach to ecology. Observations, over temporal and spatial scales well beyond standard
434 controlled experiments following mass die-offs (Levitan 1988, Linares et al. 2005), large
435 disturbance events such as ice scour (McCook & Chapman 1997), storms (Hughes 1989,
436 Underwood 1998), earthquakes (Castilla 1988), human management intervention (Hughes et
437 al. 2009) and pollution (Southward & Southward 1978) have created insight into ecological
438 processes. McCook and Chapman (1997) assessed the impacts of large scale ice scouring on
439 exposed rocky shores in Nova Scotia over a 5 year period, and showed that many
440 experimental effects noted over shorter time scales (McCook & Chapman 1991,1992) were
441 not apparent after longer periods, suggesting that much of the variation within the exposed
442 shore habitat is temporary. They also argue that the temporal dominance of *Fucus* observed
443 over mussels over a 5 year period, and no evidence of competitive superiority by mussels,
444 should lead to caution in assuming any general applicability of previously stated general
445 models of community structure based on the work of Menge and Sutherland (1976, 1987).
446 Such conclusions, based on long term observations following large scale perturbations,
447 present a cautionary note to basing environmental management advice on a theoretical
448 framework based predominantly on the short term and small scale. Underwood (1998) makes
449 the point that observations following disturbance can give rise to hypotheses regarding
450 mechanisms which can then be tested through small scale, shorter term experiments. He
451 observed variable rates of recovery in the canopy alga *Hormosira banksii* over a seven year
452 period following an extremely large storm and then used short term experimental
453 manipulations to test two alternative models: 1) that the rate of recovery was a function of the
454 severity of disturbance; 2) recovery was determined by the activities of grazers. A similar
455 combined approach was taken by Carroll and Highsmith (1996). A severe freeze in Alaska
456 coincided with a spring low tide series leading to high levels of mussel mortality, but little

effect on the mussels' principal predator *Nucella lima* which winters in the sub-tidal. A combination of observations over 3 years and experimental manipulations demonstrated that the extreme event 'set the framework for a shift in balance of the interaction' between predator and prey (Carroll & Highsmith 1996, p.130). In the presence of high numbers of *N. lima*, mussels were unable to recover over a 3 year period despite annual recruitment events.

One type of 'experiment' in coastal marine environments which by its nature tends to be long term is the implementation of Marine Protected Areas (MPA's). By totally or partially excluding the extractive activities of humans (Sciberras et al. 2015) MPA's establish a (usually) large scale experiment (Castilla 1999, Diaz et al. 2003) which can be used for understanding long-term dynamics (Castilla 2000) and has led to enormous ecological insight in coastal marine systems, particularly into the prevalence, time scale and magnitude of trophic cascades (reviewed by Castilla 1999). For example in Chile, Duran and Castilla (1989) report the results of a five year study of the consequences of exclusion of human food gatherers from the rocky intertidal and shallow sub-tidal at Las Cruces in central Chile. The cessation of human exploitation of the carnivorous muricid gastropod *Concholepas concholepas* or 'loco' led to significant enhancement of its density and consequent strong predatory control of previously dominating mussels. These cascading effects resulted in an elevation of intertidal diversity and subsequent domination by barnacles.

The majority of marine reserves or protected areas are established to protect fisheries-targeted species, including benthic organisms such as lobsters, scallops, abalone and the Chilean 'loco'. Such organisms are generally long lived relative to the majority of benthic organisms studied in perturbation experiments. Because the focus of MPA studies has often been to establish effects on these relatively long lived target species (e.g. Duran & Castilla 1989, Hoskin et al. 2011), and because the temporal scale of the press manipulation (reduction or elimination of human disturbance) is not limited by the normal drivers of ecological experimentation (e.g. researchers funds), it is not surprising that published studies of MPAs are often long term. Harvested species may show a significant lag in response (e.g. Beukers-Stewart et al. 2005), but benthic prey may also show many years to respond to enhanced predation. For example sea urchins can take many years to respond to changes in predator numbers, (e.g. 13 years in New Zealand, 7 years, in Tasmania, 15 years in Kenya; see Babcock et al. 2010 and references therein). Thus indirect effects on organisms that occur through cascading trophic interactions can take substantially longer to develop than direct effects (Babcock et al. 2010). This observation may explain why species assemblages in

older reserves often differ from recently created reserves (Micheli et al. 2004) and could lead to failure to identify trophic cascades owing to a lack of sufficient temporal scale (Shears & Babcock 2003).

The large scale, uncontrolled nature of MPA experiments can lead to considerable insight into the nature of the relationship between environmental and biotic drivers of community state when observations are conducted over sufficient time scale (Babcock et al. 2010, Fraschetti et al. 2013). Fraschetti et al. (2013, p.11) consider a long term approach is essential because *'ecological components may respond differently and over varying temporal scales'*. They used a nine year series of observations inside and outside an MPA in SE Italy to examine the role of protection in determining temporal and spatial variability. They showed that protected assemblages, with relatively long lived invertebrates and macroalgae, displayed spatial and temporal homogeneity in marked contrast to a mosaic of changing disturbed patches outside the MPA. Thus, at a local scale, protection enhanced community stability. However, there were also a number of idiosyncratic results, where the effects of protection did not match expectations. Here, it is likely that multiple processes, some derived from variation in environmental context, were acting simultaneously. Similarly Shears and Babcock (2003) demonstrated long term effects (25 years) of protection in a trophic cascade in the Leigh marine reserve in New Zealand mediated through the decline of urchins. However, urchin decline was also observed over the short term (possibly as a consequence of disease) at some unprotected sites.

Studies of the effect of protection from human disturbance are often made at a single point in time through comparison of protected and unprotected sites, often many years after establishment of the protection regime. Although suffering from difficulties in firmly ascribing observed spatial differences to protection effects (Underwood 1991), these studies are logistically relatively easy to implement and can (limitations accepted) give important long term insight. For example Mumby et al. (2006) sampled fish and benthic assemblages of Caribbean reef systems 18 years after implementation of a fishing ban. A net doubling of grazing pressure from parrot fishes in protected areas led to a four-fold reduction in macroalgal cover (Mumby et al. 2006) and consequently a two fold increase in coral recruitment (Mumby et al. 2007). Clearly such observations tell us nothing of the rate of change in fish or benthic assemblages, nor the temporal dynamic between interacting species, but does demonstrate the strong top down control exerted by key fish consumers in supporting the health of Caribbean coral reefs. As a note of caution to accepting conclusions based on

such an approach, a review of six time series studies showed that increases in target species following protection were not always as high as expected on the basis of studies with no temporal replication (Babcock et al. 2010). Thus, understanding how communities change through time following protection is likely to lead to a far greater mechanistic understanding of protection effects.

Conclusions

It is clear from our review of experimental duration over 33 years, that the majority of manipulative experiments in benthic systems, examining community or population responses of macro-organisms, are reasonably short (two years or less). It is also clear, from reviewing a range of experiments operating over a longer time scale, that a long term approach can yield considerable insight, and in many cases alter conclusions based on the same experiments conducted over shorter time scales. Thus we argue that although many short studies will be perfectly valid, in general, increasing the period over which observations are made following experimental manipulations will yield important benefits.

Limitations on the temporal extent of marine benthic experiments are numerous, and range from the logistical to considerations of motive and reward. Raffaelli and Moller (2000) showed through interviews of experimentalists that decisions regarding duration of field experiments were predominantly pragmatic. Temporal scale was often based on the time available in a studentship, or grant, or the difficulties of maintaining the experiment in a hostile environment. However, there are a number of simple, practical approaches that can be implemented to lengthen experimental duration. First and foremost when planning manipulative field experiments Diaz et al. (2003) stress the need for foresight and the acquisition of long-term funding where possible. However, extending sampling regimes may not necessarily be costly. A common approach where the temporal scale has been extended is to sample at regular intervals over the first year or so of an experiment and then sample at much lower frequencies over a number of years (e.g. Jenkins et al. 2004, Viejo 2009). Extending an experiment over multiple years generally requires some foresight in experimental establishment (e.g. use of secure, non corrosive, preferably stainless steel, markers and a mapping system of spatial replicates which can be utilised by persons other than the original experimenter), plus the motivation to revisit experimental sites over multiple

years. This can be achieved long after the original experimenter (e.g. student, post-doctoral scientist) has moved on to new pastures.

In promoting a longer term approach to experimentation it is reasonable to consider the role of reviewers and editors. Working over an appropriate temporal scale to address the question posed is an integral part of experimental design and hence should be considered as an issue ranking in importance alongside appropriate replication and independence of sampling units. Whether it currently is, or not, is debatable. That may be because the question is much less tangible. It is always likely that extending an experiment will give additional insight but the cost-benefit relationship of extended monitoring for potentially marginal gains is probably complex and difficult to pin down. However, it is clear from the review of long term studies above, that an increased mechanistic understanding of marine ecological processes can be gained from a longer term approach and understanding the longer-term consequences of impacts on benthic systems is a crucial part of determining management options in the face of global change.

Acknowledgements

This work was supported by a Brazilian FAPESP Visiting Researcher grant to the first author. Facilities were kindly supplied by the Departamento de Ciências do Mar, Universidade Federal de São Paulo - Campus Baixada Santista, Brazil. The manuscript was much improved by the comments of five anonymous reviewers.

References

- Aberg P (1992) Size based demography of the seaweed *Ascophyllum nodosum* in stochastic environments. *Ecology* 73:1488-1501
- Aberg P, Pavia H (1997) Temporal and multiple scale variation in juvenile and adult abundance of the brown alga *Ascophyllum nodosum*. *Mar Ecol Prog Ser* 158:111-119
- Araújo RM, Sousa-Pinto I, Serrão EA, Åberg P (2012) Recovery after trampling disturbance in a canopy-forming seaweed population. *Mar Biol* 159:697–707

584 Babcock RC, Shears NT, Alcala AC, Barrett NS, Edgar GJ, Lafferty KD, McClanahan TR,
 585 Russ GR (2010) Decadal trends in marine reserves reveal differential rates of change
 586 in direct and indirect effects. *Proc Natl Acad Sci USA* 107:18256-18261
 587 Barnes DKA (1996) Low levels of colonisation in Antarctica: the role of bryozoans in early
 588 community development. In: Gordon D, Smith A, Grant-Mackie A (eds) *Bryozoans in*
 589 *space and time*. National Institute of Water & Atmospheric Research Ltd. Wellington,
 590 p 19–28
 591 Benedetti-Cecchi L (2003) The importance of the variance around the mean effect size of
 592 ecological processes. *Ecology* 84:2335-2346
 593 Bertness MD, Callaway R (1994) Positive interactions in communities. *Trends Ecol Evol*
 594 9:191-193
 595 Bertness MD, Trussell GC, Ewanchuk PJ, Silliman BR (2002) Do alternate stable community
 596 states exist in the Gulf of Maine rocky intertidal zone? *Ecology* 83:3434-3448
 597 Beuchel F, Gulliksen B (2008) Temporal patterns of benthic community development in an
 598 Arctic fjord (Kongsfjorden, Svalbard): results of a 24-year manipulation study. *Polar*
 599 *Biology* 31:913-924
 600 Beukers-Stewart BD, Vause BJ, Mosley MWJ, Rossetti HL, Brand AR (2005) Benefits of
 601 closed area protection for a population of scallops. *Mar Ecol Prog Ser* 298:189–204,
 602 Broitman BR, Navarrete SA, Smith F, Gaines SD (2001) Geographic variation of
 603 southeastern Pacific intertidal communities. *Mar Ecol Prog Ser* 224:21-34
 604 Carroll ML, Highsmith RC (1996) Role of catastrophic disturbance in mediating *Nucella-*
 605 *Mytilus* interactions in the Alaskan rocky intertidal. *Mar Ecol Prog Ser* 138:125-133
 606 Castilla JC (1988) Earthquake-caused coastal uplift and its effects on rocky intertidal kelp
 607 communities. *Science* 242:440–443
 608 Castilla JC (1999) Coastal marine communities: trends and perspectives from human-
 609 exclusion experiments. *Trends Ecol Evol* 14:280-283
 610 Castilla JC (2000) Roles of experimental marine ecology in coastal management and
 611 conservation. *J Exp Mar Biol Ecol* 250:3-21
 612 Castilla JC, Bustamante R (1989) Human exclusion from rocky intertidal of Las-Cruces,
 613 central Chile - effects on *Durvillaea-antarctica* (Phaeophyta, durvilleales). *Mar Ecol*
 614 *Prog Ser* 50:203-214
 615 Cervin G, Aberg P, Jenkins SR (2005) Small-scale disturbance in a stable canopy dominated
 616 community: implications for macroalgal recruitment and growth. *Mar Ecol Prog Ser*
 617 305:31-40
 618 Chapman MG (2013) Constructing replacement habitat for specialist and generalist molluscs-
 619 the effect of patch size. *Mar Ecol Prog Ser* 473:201-214
 620 Coleman RA, Underwood AJ, Benedetti-Cecchi L, Aberg P, Arenas F, Arrontes J, Castro J,
 621 Hartnoll RG, Jenkins SR, Paula J, Della Santina P, Hawkins SJ (2006) A continental
 622 scale evaluation of the role of limpet grazing on rocky shores. *Oecologia* 147:556-564
 623 Connell JH (1961) The influence of interspecific competition and other factors on the
 624 distribution of the barnacle *Chthamalus stellatus*. *Ecology* 42:710-723
 625 Connell JH (1972). Community interactions on marine rocky intertidal shores. *Annu Rev Ecol*
 626 *Syst* 3: 169-192
 627 Connell JH, Slayter RO (1977) Mechanisms of succession in natural communities and their
 628 role in community stability and organization. *Am Nat* 111:1119-1144

629 Connell JH, Sousa WP (1983) On the evidence needed to judge ecological stability or
 630 persistence *Am Nat* 121:789-824
 631 Connell SD (2005) Assembly and maintenance of subtidal habitat heterogeneity: synergistic
 632 effects of light penetration and sedimentation. *Mar Ecol Prog Ser* 289:53-61
 633 Connell SD, Foster MS, Airoidi L (2014) What are algal turfs? Toward a better
 634 understanding of turfs. *Mar Ecol Prog Ser* 495:299-307
 635 Connolly SR, Roughgarden J (1998) A latitudinal gradient in northeast Pacific intertidal
 636 community structure: Evidence for an oceanographically based synthesis of marine
 637 community theory. *Am Nat* 151:311-326
 638 Crowe TP, Bracken MES, O'Connor NE (2012) Reality check: issues of scale and abstraction
 639 in biodiversity research and potential solutions. In: Solan M, Aspden R, Paterson D
 640 (eds) *Marine biodiversity and ecosystem functioning: frameworks, methodologies and*
 641 *integration*. Oxford University Press Oxford
 642 Davis AR, Ward DW (2009) Establishment and persistence of species-rich patches in a
 643 species-poor landscape: role of a structure-forming sub-tidal barnacle. *Mar Ecol Prog*
 644 *Ser* 380:187-198
 645 Dayton PK (1975) Experimental evaluation of ecological dominance in a rocky intertidal
 646 algal community. *Ecol Monogr* 45:137-159
 647 Dayton PK (1989) Inter-decadal variation in an Antarctic sponge and its predators from
 648 oceanographic climate shifts. *Science* 245:1484–1486
 649 Dayton PK, Oliver JS (1977) Antarctic soft-bottom benthos in oligotrophic and eutrophic
 650 environments. *Science* 197:55–58
 651 Diaz S, Symstad AJ, Chapin FS, Wardle DA, Huenneke LF (2003) Functional diversity
 652 revealed by removal experiments. *Trends Ecol Evol* 18:140-146
 653 Duffy JE, Stachowicz JJ, Bruno JF (2012) Multi-trophic biodiversity and the responses of
 654 marine ecosystems to global change In: Solan M, Aspden R, Paterson D (eds) *Marine*
 655 *biodiversity and ecosystem functioning: frameworks, methodologies and integration*.
 656 Oxford University Press, Oxford
 657 Duran LR, Castilla JC (1989) Variation and persistence of the middle rocky intertidal
 658 community of central Chile, with and without human harvesting. *Mar Biol* 103:555-
 659 562
 660 Dye AH (1998) Community-level analyses of long-term changes in rocky littoral fauna from
 661 South Africa. *Mar Ecol Prog Ser* 164:47-57
 662 Ellis J, Schneider DC (2008) Spatial and temporal scaling in benthic ecology. *J Exp Mar Biol*
 663 *Ecol* 366:92-98
 664 Farrell TM (1988) Community stability: effects of limpet removal and reintroduction in a
 665 rocky intertidal community. *Oecologia* 75:190-197
 666 Feldman KL, Armstrong DA, Eggleston DB, Dumbauld BR (1997) Effects of substrate
 667 selection and post-settlement survival on recruitment success of the thalassinidean
 668 shrimp *Neotrypaea californiensis* to inter-tidal shell and mud habitats. *Mar Ecol Prog*
 669 *Ser* 150:121-136
 670 Fraschetti S, Guarnieri G, Bevilacqua S, Terlizzi A, Boero F (2013) Protection enhances
 671 community and habitat stability: evidence from a Mediterranean marine protected
 672 area. *Plos ONE* 8(12): e81838. doi:10.1371/journal.pone.0081838
 673 Goodsell PJ, Connell SD (2005) Historical configuration of habitat influences the effects of
 674 disturbance on mobile invertebrates. *Mar Ecol Prog Ser* 299:79-87

675 Grant DL, Clarke PJ, Allaway WG (1993) The response of grey mangrove (*Avicennia marina*
676 (Forsk.) Vierh.) seedlings to spills of crude oil. *J Exp Mar Biol Ecol* 171: 273-295.

677 Green DS, Crowe TP (2013) Physical and biological effects of introduced oysters on
678 biodiversity in an intertidal boulder field. *Mar Ecol Prog Ser* 482:119-132

679 Holmes NJ, Harriott VJ, Banks SA (1997) Latitudinal variation in patterns of colonisation of
680 cryptic calcareous marine organisms. *Mar Ecol Prog Ser* 155:103-113

681 Hooper DU, Chapin FS, Ewel JJ, Hector A, Inchausti P, Lavorel S, Lawton JH, Lodge DM,
682 Loreau M, Naeem S, Schmid B, Setälä H, Symstad AJ, Vandermeer J, Wardle DA
683 (2005) Effects of biodiversity on ecosystem functioning: A consensus of current
684 knowledge. *Ecol Monogr* 75:3-35

685 Hoskin MG, Coleman RA, von Carlshausen E, Davis CM (2011) Variable population
686 responses by large decapod crustaceans to the establishment of a temperate marine no-
687 take zone. *Can J Fish Aquat Sci* 68:185-200

688 Hughes TP Community structure and diversity of coral reefs: the role of history. *Ecology* 70:
689 275-279

690 Hughes RG, Fletcher PW, Hardy MJ (2009) Successional development of saltmarsh in two
691 managed realignment areas in SE England, and prospects for saltmarsh restoration.
692 *Mar Ecol Prog Ser* 384:13-22

693 Incera M, Bertocci I, Benedetti-Cecchi L (2010) Effects of mean intensity and temporal
694 variability of disturbance on the invasion of *Caulerpa racemosa* var. *cylindracea*
695 (*Caulerpales*) in rock pools. *Biol Inv* 12:501-514

696 Incera M, Olabarria C, Troncoso JS, Lopez J (2009) Response of the invader *Sargassum*
697 *muticum* to variability in nutrient supply. *Mar Ecol Prog Ser* 377:91-101

698 Ingolfsson A, Hawkins SJ (2008) Slow recovery from disturbance: a 20 year study of
699 *Ascophyllum* canopy clearances. *J Mar Biol Assoc UK* 88:689-691

700 Jacobs R (1980) Effects of the Amoco-Cadiz oil-spill on the seagrass community at Roscoff
701 with special reference to the benthic infauna. *Mar Ecol Prog Ser* 2:207-212

702 Jenkins SR (2009) Settlement and recruitment In: M W (ed) *Marine Hard Bottom*
703 *Communities: Patterns, Dynamics, Diversity and Change*. Springer, Verlag

704 Jenkins SR, Arenas F, Arrontes J, Bussell J, Castro J, Coleman RA, Hawkins SJ, Kay S,
705 Martinez B, Oliveros J, Roberts MF, Sousa S, Thompson RC, Hartnoll RG (2001)
706 European-scale analysis of seasonal variability in limpet grazing activity and
707 microalgal abundance. *Mar Ecol Prog Ser* 211:193-203

708 Jenkins SR, Coleman RA, Della Santina P, Hawkins SJ, Burrows MT, Hartnoll RG (2005)
709 Regional scale differences in the determinism of grazing effects in the rocky intertidal.
710 *Mar Ecol Prog Ser* 287:77-86

711 Jenkins SR, Hawkins SJ, Norton TA (1999a) Direct and indirect effects of a macroalgal
712 canopy and limpet grazing in structuring a sheltered inter-tidal community. *Mar Ecol*
713 *Prog Ser* 188:81-92

714 Jenkins SR, Norton TA, Hawkins SJ (1999b) Interactions between canopy forming algae in
715 the eulittoral zone of sheltered rocky shores on the Isle of Man. *J Mar Biol Assoc UK*
716 79:341-349

717 Jenkins SR, Norton TA, Hawkins SJ (2004) Long term effects of *Ascophyllum nodosum*
718 canopy removal on mid shore community structure. *J Mar Biol Assoc UK* 84:327-329

719 Jewett EB, Hines AH, Ruiz GM (2005) Epifaunal disturbance by periodic low levels of
720 dissolved oxygen: native vs. invasive species response. *Mar Ecol Prog Ser* 304:31-44

721 Kaehler S, Williams GA (1997) Do factors influencing recruitment ultimately determine the
722 distribution and abundance of encrusting algae on seasonal tropical shores? *Mar Ecol*
723 *Prog Ser* 156:87-96

724 Keough MJ, Downes BJ (1982) Recruitment of marine-invertebrates - the role of active larval
725 choices and early mortality. *Oecologia* 54:348-352

726 Kilar JA, McLachlan J (1989) Effects of wave exposure on the community structure of a
727 plant-dominated, fringing-reef platform - intermediate disturbance and disturbance-
728 mediated competition. *Mar Ecol Prog Ser* 54:265-276

729 Konar B (2007) Recolonization of a high latitude hard-bottom nearshore community. *Polar*
730 *Biology* 30:663-667

731 Leonard GH (1999) Positive and negative effects of intertidal algal canopies on recruitment
732 and survival of barnacles. *Mar Ecol Prog Ser* 178:241-249

733 Levin S (1992) The problem of pattern and scale in ecology. *Ecology* 72:1943-1967

734 Levitan DR (1988) Algal-urchin biomass responses following mass mortality of *Diadema*
735 *antillarum* Philippi at Saint John, U.S. Virgin Islands. *J Exp Mar Biol Ecol*
736 119:167-178

737 Lewontin RC (1969) The meaning of stability. In: Woodwell G, Smith H (eds) *Diversity and*
738 *stability in ecological systems*. Brookhaven National Laboratory, Upton, NY., p 12-24

739 Linares C, Coma R, Diaz D, Zabala M, Hereu B, Dantart L (2005) Immediate and delayed
740 effects of a mass mortality event on gorgonian population dynamics and benthic
741 community structure in the NW Mediterranean Sea. *Mar Ecol Prog Ser* 305:127-137

742 Lindegarth M, Aberg P, Cervin G, Nilsson PG (2001) Effects of grazing on the structure of
743 mid-shore, inter-tidal assemblages on moderately exposed rocky shores of the
744 Swedish west coast. *Mar Ecol Prog Ser* 212:29-38

745 Lubchenco J, Menge BA (1978) Community development and persistence in a low rocky
746 intertidal zone. *Ecol Monogr* 48:67-94

747 Maggi E, Bertocci I, Vaselli S, Benedetti-Cecchi L (2009) Effects of changes in number,
748 identity and abundance of habitat-forming species on assemblages of rocky seashores.
749 *Mar Ecol Prog Ser* 381:39-49

750 Maggi E, Bulleri F, Bertocci I, Benedetti-Cecchi L (2012) Competitive ability of macroalgal
751 canopies overwhelms the effects of variable regimes of disturbance. *Mar Ecol Prog*
752 *Ser* 465:99-109

753 McCook, LJ Chapman ARO (1991) Community succession following massive ice scour on
754 an exposed rocky shore : effects of *Fucus* canopy algae and of mussels during late
755 succession. *J Exp Mar Biol Ecol* 154:137-169

756 McCook LJ, Chapman ARO (1992) Vegetative regeneration of *Fucus* rockweed canopy as a
757 mechanism of secondary succession on an exposed rocky shore. *Bot Mar* 35:35-

758 McCook LJ, Chapman ARO (1997) Patterns and variations in natural succession following
759 massive ice-scour of a rocky intertidal seashore. *J Exp Mar Biol Ecol* 214:121-147

760 Menge BA (1997) Detection of direct versus indirect effects: Were experiments long enough?
761 *Am Nat* 149:801-823

762 Menge BA, Daley BA, Wheeler PA, Dahlhoff E, Sanford E, Strub PT (1997) Benthic-pelagic
763 links and rocky intertidal communities: Bottom-up effects on top-down control? *Proc*
764 *Natl Acad Sci USA* 94:14530-14535

765 Menge BA, Lubchenco J, Bracken MES, Chan F, Foley MM, Freidenburg TL, Gaines SD,
766 Hudson G, Krenz C, Leslie H, Menge DNL, Russell R, Webster MS (2003) Coastal

767 oceanography sets the pace of rocky intertidal community dynamics. *Proc Natl Acad*
768 *Sci USA* 100:12229-12234

769 Menge BA, Sanford E, Daley BA, Freidenburg TL, Hudson G, Lubchenco J (2002) Inter-
770 hemispheric comparison of bottom-up effects on community structure: Insights
771 revealed using the comparative-experimental approach. *Ecol Res* 17:1-16

772 Menge BA, Sutherland JP (1976) Species diversity gradients: synthesis of the roles of
773 predation, competition and temporal heterogeneity. *Am Nat* 110:351-369

774 Menge BA, Sutherland JP (1987) Community regulation: Variation in disturbance,
775 competition and predation in relation to environmental stress and recruitment. *Am Nat*
776 130:730-757

777 Methratta ET, Petraitis PS (2008) Propagation of scale-dependent effects from recruits to
778 adults in barnacles and seaweeds. *Ecology* 89:3128-3137

779 Micheli F, Halpern BS, Botsford LW, Warner RR (2004) Trajectories and correlates of
780 community change in no-take marine reserves. *Ecol Appl* 14:1709–1723

781 Miller MW, Valdivia A, Kramer KL, Mason B, Williams DE, Johnston L (2009) Alternate
782 benthic assemblages on reef restoration structures and cascading effects on coral
783 settlement. *Mar Ecol Prog Ser* 387:147-156

784 Minchinton TE, Fels KJ (2013) Sediment disturbance associated with trampling by humans
785 alters species assemblages on a rocky intertidal seashore. *Mar Ecol Prog Ser* 472:129-
786 140

787 Mumby PJ, Dahlgren CP, Harborne AR, Kappel CV, Micheli F, Brumbaugh DR, Holmes KE,
788 Mendes JM, Broad K, Sanchirico JN, Buch K, Box S, Stoffle RW, Gill AB (2006)
789 Fishing, trophic cascades, and the process of grazing on coral reefs. *Science* 311:98-
790 101

791 Mumby PJ, Harborne AR, Williams J, Kappel CV, Brumbaugh DR, Micheli F, Holmes KE,
792 Dahlgren CP, Paris CB, Blackwell PG (2007) Trophic cascade facilitates coral
793 recruitment in a marine reserve. *Proc Natl Acad Sci USA* 104:8362-8367

794 Naeem S (2008) Advancing realism in biodiversity research. *Trends in Ecology and*
795 *Evolution* 23:414-416

796 Nelson WG (1981) Experimental studies of decapod and fish predation on seagrass
797 macrobenthos. *Mar Ecol Prog Ser* 5:141-149

798 O'Connor NE, Crowe TP (2005) Biodiversity loss and ecosystem functioning: Distinguishing
799 between number and identity of species. *Ecology* 86:1783-1796

800 Oliveira JP, Sousa-Pinto I, Weber GM, Bertocci I (2014) Interplay of experimental harvesting
801 and climate-related disturbance on benthic assemblages of rocky seashores. *Mar Ecol*
802 *Prog Ser* 495:131-142

803 Paine RT (1966) Food web complexity and species diversity. *Am Nat* 100:65-75

804 Paine, R. (1977). Controlled manipulations in the marine intertidal zone and their
805 contributions to ecological theory. In: Goulden CE (ed) *Changing scenes in natural*
806 *sciences 1776-1976*. Academy of Natural Sciences, Philadelphia, PA, 245-270.

807 Paine RT (1994) Marine rocky shores and community ecology: an experimentalist's
808 perspective. In: Kinne O. (ed.) *Excellence in Ecology 4*. Ecology Institute, D-21385
809 Oldendorf/Luhe, Germany

810 Paine RT, Castillo JC, Cancino J (1985) Perturbation and recovery patterns of starfish-
811 dominated intertidal assemblages in Chile, New Zealand, and Washington State. *Am*
812 *Nat* 125:679-691

813 Peterson CH (1984) Does a rigorous criterion for environmental identity preclude the
814 existence of multiple stable points? *Am Nat* 124:127-133

815 Petraitis PS, Dudgeon SR (2004) Detection of alternative stable states in marine communities.
816 *J Exp Mar Biol Ecol* 300:343- 371

817 Petraitis PS, Dudgeon SR (2005) Divergent succession and implications for alternative states
818 on rocky intertidal shores. *J Exp Mar Biol Ecol* 326:14 – 26

819 Petraitis PS, Latham RE (1999) The importance of scale in testing the origins of alternative
820 community states. *Ecology* 80: 429 - 442

821 Petraitis PS, Methratta ET, Rhile EC, Vidargas NA, Dudgeon SR (2009) Experimental
822 confirmation of multiple community states in a marine ecosystem. *Oecologia* 161:139-
823 148

824 Raffaelli DG (2006) Biodiversity and ecosystem functioning: issues of scale and trophic
825 complexity. *Mar Ecol Prog Ser* 311:285-294

826 Raffaelli DG, Hawkins SJ (1996) Intertidal ecology. Chapman and Hall, London

827 Raffaelli DG, Moller H (2000) Manipulative experiments in animal ecology-Do they promise
828 more than they can deliver? *Adv Ecol Res* 30:299-330

829 Reinhardt JF, Whitlatch RB, Osman RW (2013) Effects of temperature on the recruitment
830 phenology and niche overlap of shallow epifaunal assemblages in southern New
831 England. *Mar Ecol Prog Ser* 489:61-74

832 Ruiz JM, Romero J (2001) Effects of in situ experimental shading on the Mediterranean
833 seagrass *Posidonia oceanica*. *Mar Ecol Prog Ser* 215:107-120

834 Russell BD, Connell SD (2005) A novel interaction between nutrients and grazers alters
835 relative dominance of marine habitats. *Mar Ecol Prog Ser* 289:5-11

836 Sanford E (1999) Regulation of keystone predation by small changes in ocean temperature.
837 *Science* 283:2095-2097

838 Scheffer M, Carpenter S, Foley JA, Folke C, Walker B (2001) Catastrophic shifts in
839 ecosystems. *Nature* 413:591 - 596

840 Schiel DR, Lilley SA (2011) Impacts and negative feedbacks in community recovery over
841 eight years following removal of habitat-forming macroalgae. *J Exp Mar Biol Ecol*
842 407:108-115

843 Schneider DC (2001) The rise of the concept of scale in ecology. *Bioscience* 51:545-553

844 Sciberras M, Jenkins SR, Kaiser MJ, Hawkins SJ, Pullin AS (2015) Evaluating the relative
845 conservation value of partially and fully protected marine reserves. *Fish and Fisheries*
846 16: 58–77

847 Shears NT, Babcock RC (2003) Continuing trophic cascade effects after 25 years of no take
848 marine reserve protection. *Mar Ecol Prog Ser* 246:1-16

849 Simenstad CA, Estes JA, Kenyon KW (1978) Aleuts, sea otters, and alternate stable state
850 communities. *Science* 200:403-411

851 Southward AJ, Southward EC (1978) Recolonization of rocky shores in Cornwall after use of
852 toxic dispersants to clean up the Torrey Canyon spill. *J Fish Res Board Can* 35:682-
853 706

854 Spivak AC, Canuel EA, Duffy JE, Douglass JG, Richardson JP (2009) Epifaunal community
855 composition and nutrient addition alter sediment organic matter composition in a
856 natural eelgrass *Zostera marina* bed: a field experiment. *Mar Ecol Prog Ser* 376:55-67

- Stachowicz JJ, Bracken MES, Graham M (2008b) Complementarity in marine biodiversity manipulations: reconciling divergent evidence from field and mesocosm experiments. *Proc Natl Acad Sci* 105:18842-18847
- Stachowicz JJ, Graham M, Bracken MES, Szoboszlai AI (2008a) Diversity enhances cover and stability of seaweed assemblages: the role of heterogeneity and time. *Ecology* 89:3008–3019
- Sutherland JP (1990) Perturbations, resistance and alternative views of the existence of multiple stable points in nature. *Am Nat* 136:270-275
- Sutherland JP, Karlson RH (1977) Development and stability of fouling community at Beaufort, North-Carolina. *Ecol Monogr* 47:425-446
- Tait LW, Schiel DR (2011) Legacy effects of canopy disturbance on ecosystem functioning in macroalgal assemblages. *PloS ONE* DOI: 10.1371/journal.pone.0026986
- Tegner MJ, Dayton PK, Edwards PB, Riser KL (1997) Large-scale, low-frequency oceanographic effects on kelp forest succession: A tale of two cohorts. *Mar Ecol Prog Ser* 146:117-134
- Underwood AJ (1991) Beyond BACI: Experimental designs for detecting human environmental impacts on temporal variations in natural populations. *Aust J Mar Freshwat Res* 42:569-587
- Underwood AJ (1998) Grazing and disturbance: an experimental analysis of patchiness in recovery from a severe storm by the intertidal alga *Hormosira banksii* on rocky shores in New South Wales. *J Exp Mar Biol Ecol* 231:291-306
- Underwood AJ (2000) Experimental ecology of rocky intertidal habitats: what are we learning? *J Exp Mar Biol Ecol* 250:51-76
- Valdivia N, Heidemann A, Thiel M, Molis M, Wahl M (2005) Effects of disturbance on the diversity of hard-bottom macrobenthic communities on the coast of Chile. *Mar Ecol Prog Ser* 299:45-54
- Van De Koppel J, Herman PMJ, Thoolen P, Heip CHR (2001) Do alternate stable states occur in natural ecosystems? Evidence from a tidal flat. *Ecology* 82:3449 - 3461
- Viejo RM (2009) Resilience in intertidal rocky shore assemblages across the stress gradient created by emersion times. *Mar Ecol Prog Ser* 390:55-65
- Wiens JA (1989) Spatial scaling in ecology *Func Ecol* 3:385-397
- Yodzis P (1988) The indeterminacy of ecological interactions as perceived through perturbation experiments *Ecology* 69:508-515




Figure 1 Frequency histogram indicating the distribution of experimental study duration in (A) marine journals and (B) ecological journals over the period 1980 to 2013.





Figure 2 Box plot showing the duration of manipulative experiments over the nine periods examined and for all periods combined in (A) marine journals and (B) ecological journals. Boxes indicate the median (horizontal line) and the 25th and 75th percentiles; whiskers indicate the 10th and 90th percentile and black dots the outliers. Number of independent experiments included in the analysis indicated in brackets beneath each year.



915

916

917

918

919

920

921

922

Figure 3 Box plots showing median study duration (and 10th, 25th, 75th and 90th percentiles) for comparisons among A) press (n=219) and pulse (n=133) studies and B) intertidal (n= 175) and sub-tidal (n=177) studies